

HEPATITIS C VIRUS SPECIALTY CARE PROGRAM

Phone: **844-422-6400** Fax: **888-850-4018**

KLOUDSCRIPT Community Led Specialty Pharmacy Care

v9.4_080817

©2017 KloudScript, Inc. - All rights reserved.

PATIENT INFORMATI			BER INFORMATION:	
Name:				
Address: City:	State: Zin:	Address	State: Z	in:
•	-	-		•
Phone:			Fax:	
			DEA:	
DOB: Gender: O	_			
Height: Weight:	Allergies:	Office Contact:	Phone:	
3 STATEMENT OF MEI	DICAL NECESSITY: (Plea	ase Attach All Medical	Documentation)	
	Diagnostic Information		Labs	
Date of Diagnosis: ICD-	-10: Other:		ALT: HGB:	
Genotype: Subtype: _	Subtype: Q80K: Positive Negative (fit Status: Naïve Partial Responder Non-responder Null-resp		AST: HCV RN	
			PLT: SrCr:	
Duration of Previous Therapy:Cirrhosis: ☐ No ☐ Yes If Yes: ☐ Co				
History of Liver Biopsy? No Yes			NS5A Resistance Assay:	Date:
☐ Fibrosure or ☐ Fibroscan: Results	-		Medication List and Contra	indications
Extra-Hepatic Manifestations: Asc			☐ Attach Medication List	
Other:			☐ Yes ☐ No ☐ Is the patient interferon ineligible? ☐ No ☐ Yes	
HBsAg and anti-HBc Test: ☐ Positive	e 🛘 Negative Date:		l '	
If Prior Authorization is Denied: ☐ Automatically Draft Appeal for Re	Donal Droferred Formulars	Allamatina	☐ Anxiety ☐ Depression ☐ Pulmona	
Automatically Draft Appeal for ne	Wiew 🔲 Sena Preierrea Formalary	/ Alternatives	Renal Insufficiency Other:	
4 INJECTION TRAININ	G: O To Be Administered by Pharmacist	t O Pharmacist to Provide Trainin	g O Patient Trained in MD Office O Manuf	facturer Nurse Suppo
5 PICK UP OR DELIVE	RY: O Delivery to Patient's	Home O Delivery to I	Physician's Office O Pharmacy t	o Coordinate
PRESCRIPTION: Duration of Patient Name:	of Therapy: 🛘 8 Weeks 🖵 12 W			
			tient's Date of Birth:	
Medication	Dosage & Strength		Direction	QTY Refil
	Dosage & Strength □ 30mg Tablets			QTY Refil
		☐ Take 30mg daily with or w with or without Ribavirin	Direction	
Medication	□ 30mg Tablets	☐ Take 30mg daily with or w with or without Ribavirin ☐ Take 60mg daily with or w with or without Ribavirin	Direction vithout food in combination with Sovaldi®	28
Medication	□ 30mg Tablets □ 60mg Tablets	☐ Take 30mg daily with or w with or without Ribavirin☐ Take 60mg daily with or w with or without Ribavirin☐ Take 90mg daily with or w	Direction //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi®	28
Medication □ DAKLINZA [™]	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets	☐ Take 30mg daily with or w with or without Ribavirin☐ Take 60mg daily with or w with or without Ribavirin☐ Take 90mg daily with or w with or without Ribavirin☐	Direction //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® or without food	28 28 28
Medication □ DAKLINZA [™] □ EPCLUSA [®] □ HARVONI [®] □ MAVYRET [™]	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take one tablet daily with	Direction //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® or without food or without food	28 28 28 28 28
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets □ 90/400mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with	Direction without food in combination with Sovaldi® without food in combination with Sovaldi® without food in combination with Sovaldi® or without food or without food nce daily with food	28 28 28 28 28 28
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI®	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets □ 90/400mg Tablets □ 100/40mg Tablet □ 150mg Capsules □ 400mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take three tablets orally o □ Take one 150mg capsule □ Take one 400mg tablet or	Direction //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® or without food or without food or without food nce daily with food orally once a day ally once a day	28 28 28 28 28 28 1 Carton 28 28
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets □ 90/400mg Tablets □ 100/40mg Tablet □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one tablets orally or □ Take one 150mg capsule □ Take one 400mg tablet or □ Take two tablets once daily	Direction //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® or without food or without food or without food nce daily with food orally once a day ally once a day ly in the morning with a meal	28 28 28 28 28 1 Carton 28 28 28
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI®	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets □ 90/400mg Tablets □ 100/40mg Tablet □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take three tablets orally o □ Take one 150mg capsule □ Take one 400mg tablet or. □ Take two tablets once dai □ Take three tablets in the m	Direction //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® //ithout food in combination with Sovaldi® or without food or without food or without food oredaily with food orally once a day ally once a day ly in the morning with a meal norning and one tablet in the evening	28 28 28 28 28 28 1 Carton 28 28
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets □ 90/400mg Tablets □ 100/40mg Tablet □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one tablets orally or □ Take one 150mg capsule □ Take one 400mg tablet or □ Take two tablets once daily	Direction without food in combination with Sovaldi® without food in combination with Sovaldi® without food in combination with Sovaldi® or without food or without food or without food or without food orally with food orally once a day ally once a day ly in the morning with a meal morning and one tablet in the evening on the daily dose pack	28 28 28 28 28 1 Carton 28 28 28
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets □ 90/400mg Tablets □ 100/40mg Tablet □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take three tablets orally o □ Take one 150mg capsule □ Take one 400mg tablet or □ Take two tablets once dai □ Take three tablets in the m with a meal, as directed o	Direction //ithout food in combination with Sovaldi® or without food or without food ore daily with food orally once a day ally once a day ly in the morning with a meal norning and one tablet in the evening in the daily dose pack ally with food	28 28 28 28 28 28 1 Carton 28 28 56 1 Pack
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VOSEVI TM	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets □ 90/400mg Tablets □ 100/40mg Tablets □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one tablets orally o □ Take one 400mg tablet or. □ Take two tablets once dai □ Take three tablets in the m with a meal, as directed o □ Take one tablet orally onc. □ Take one tablet orally onc.	Direction //ithout food in combination with Sovaldi® or without food or without food orally with food orally once a day ally once a day ly in the morning with a meal norning and one tablet in the evening in the daily dose pack aily with food de daily with food morning/400mg tablet every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VIEKIRA XR TM □ VOSEVI TM □ MODERIBA Dose Pack TM	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets □ 90/400mg Tablets □ 100/40mg Tablet □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day □ 800mg per day	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take 90mg daily with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take three tablets orally or Take one 400mg tablet or Take two tablets once dai □ Take three tablets in the m with a meal, as directed or Take three tablets once dai □ Take one tablet orally once Take 200mg tablet every make 200mg tablet every make 400mg tablet every make 50mg tablet	Direction //ithout food in combination with Sovaldi® or without food or without food orce daily with food orally once a day ally once a day ly in the morning with a meal norning and one tablet in the evening in the daily dose pack aily with food e daily with food morning/400mg tablet every evening morning/400mg tablet every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VOSEVI TM	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 400/100mg Tablets □ 90/400mg Tablets □ 100/40mg Tablets □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one tablets orally o □ Take one 150mg capsule □ Take one 400mg tablet or □ Take two tablets once dai □ Take three tablets in the m with a meal, as directed o □ Take three tablets once dai □ Take one tablet orally onc □ Take 200mg tablet every r □ Take 400mg tablet every r □ Take 600mg tablet every r	Direction //ithout food in combination with Sovaldi® or without food or without food orally with food orally once a day ally once a day ly in the morning with a meal norning and one tablet in the evening in the daily dose pack aily with food de daily with food morning/400mg tablet every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VOSEVI TM □ VOSEVI TM □ MODERIBA Dose Pack TM □ RIBASPHERE RibaPack®	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 90/400mg Tablets □ 90/400mg Tablets □ 100/40mg Tablets □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day □ 800mg per day □ 1000mg per day □ 1200mg per day	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take 90mg daily with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one 150mg capsule □ Take one 400mg tablet or □ Take two tablets once dai □ Take three tablets in the m with a meal, as directed o □ Take three tablets once dai □ Take one tablet or ally once □ Take 200mg tablet every more □ Take 600mg tablet every more □ Take 600mg tablet every more □ Take 10mg tablet e	Direction //ithout food in combination with Sovaldi® or without food or without food or without food orally once a day ally once a day ly in the morning with a meal norning and one tablet in the evening on the daily dose pack aily with food de daily with food morning/400mg tablet every evening morning/400mg tablet every evening morning/600mg tablet every evening morning/600mg tablet every evening //capsules every morning and,	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VIEKIRA XR TM □ VOSEVI TM □ MODERIBA Dose Pack TM □ RIBASPHERE RibaPack®	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 90/400mg Tablets □ 90/400mg Tablets □ 100/40mg Tablet □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day □ 800mg per day □ 1000mg per day □ 1200mg per day	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take 90mg daily with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one 150mg capsule □ Take one 400mg tablet or □ Take two tablets once dai □ Take three tablets in the m with a meal, as directed o □ Take three tablets once dai □ Take 0ne tablet or ally once □ Take 200mg tablet every more □ Take 600mg tablet every more □	Direction //ithout food in combination with Sovaldi® or without food or without food or without food orally once a day ally once a day ly in the morning with a meal norning and one tablet in the evening on the daily dose pack aily with food de daily with food morning/400mg tablet every evening morning/400mg tablet every evening morning/600mg tablet every evening morning/600mg tablet every evening //capsules every morning and,	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VOSEVI TM □ VOSEVI TM □ MODERIBA Dose Pack TM □ RIBASPHERE RibaPack®	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 90/400mg Tablets □ 90/400mg Tablets □ 100/40mg Tablets □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day □ 800mg per day □ 1000mg per day □ 1200mg per day	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take 90mg daily with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one 150mg capsule □ Take one 400mg tablet or □ Take two tablets once dai □ Take three tablets in the m with a meal, as directed o □ Take three tablets once dai □ Take one tablet or ally once □ Take 200mg tablet every more □ Take 600mg tablet every more □ Take 600mg tablet every more □ Take 10mg tablet e	Direction ithout food in combination with Sovaldi® or without food or without food oredaily with food orally once a day ally once a day ly in the morning with a meal morning and one tablet in the evening on the daily dose pack aily with food the daily with food morning/400mg tablet every evening morning/400mg tablet every evening morning/400mg tablet every evening morning/600mg tablet every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VOSEVI TM □ VOSEVI TM □ MODERIBA Dose Pack TM □ RIBASPHERE RibaPack®	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 90/400mg Tablets □ 90/400mg Tablets □ 100/40mg Tablet □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day □ 800mg per day □ 1200mg per day □ 1200mg per day □ 1200mg Tablets □ 200mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take 90mg daily with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take three tablets orally or Take one 400mg tablet or Take two tablets once dailunced and Take three tablets in the m with a meal, as directed or Take three tablets once dailunced Take 200mg tablet every more take 400mg tablet every more take 600mg tablet every more tablets/	Direction //ithout food in combination with Sovaldi® or without food or without food oredaily with food orally once a day ally once a day ly in the morning with a meal morning and one tablet in the evening or the daily dose pack aily with food morning/400mg tablet every evening morning/400mg tablet every evening morning/400mg tablet every evening morning/600mg tablet every evening //capsules every morning and //capsules every evening y with or without food	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VOSEVI TM □ VOSEVI TM □ MODERIBA Dose Pack TM □ RIBASPHERE RibaPack® □ MODERIBA TM □ RIBASPHERE® □ RIBAVIRIN □ XIFAXAN®	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 90/400mg Tablets □ 100/40mg Tablets □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day □ 800mg per day □ 1000mg per day □ 1200mg per day □ 1200mg per day □ 1200mg Tablets □ 200mg Tablets □ 200mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take 90mg daily with or with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take three tablets orally or Take one 400mg tablet or Take two tablets once dailung the most of Take three tablets in the m with a meal, as directed or Take three tablets once dailung Take one tablet orally once Take 400mg tablet every most of Take 400mg tablet every most of Take 600mg tablet every most of Take 10mg tablet every most of Take 1	Direction //ithout food in combination with Sovaldi® or without food or without food oredaily with food orally once a day ally once a day ly in the morning with a meal morning and one tablet in the evening or the daily dose pack aily with food morning/400mg tablet every evening morning/400mg tablet every evening morning/400mg tablet every evening morning/600mg tablet every evening //capsules every morning and //capsules every evening y with or without food	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VIEKIRA XR TM □ VOSEVI TM □ MODERIBA Dose Pack TM □ RIBASPHERE RibaPack® □ MODERIBA TM □ RIBASPHERE® □ RIBAVIRIN □ XIFAXAN® □ ZEPATIER TM □	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 90/400mg Tablets □ 90/400mg Tablets □ 100/40mg Tablets □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day □ 800mg per day □ 1000mg per day □ 1200mg per day □ 1200mg Tablets □ 200mg Tablets □ 200mg Tablets □ 550mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take 90mg daily with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one 150mg capsule □ Take one 400mg tablet or Take two tablets once dai □ Take three tablets once dai □ Take three tablets once dai □ Take one tablet orally once Take 200mg tablet every round Take 400mg tablet every round Take 600mg tablet every round Take 600mg tablet every round Take 10mg tabl	Direction //ithout food in combination with Sovaldi® or without food or without food oredaily with food orally once a day ally once a day ly in the morning with a meal morning and one tablet in the evening on the daily dose pack aily with food morning/400mg tablet every evening morning/400mg tablet every evening morning/400mg tablet every evening morning/600mg tablet every evening //capsules every morning and, //capsules every evening y with or without food or without food	28 28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VIEKIRA XR TM □ VOSEVI TM □ MODERIBA Dose Pack TM □ RIBASPHERE RibaPack® □ MODERIBA TM □ RIBASPHERE® □ RIBAVIRIN □ XIFAXAN® □ ZEPATIER TM □ PRESCRIBER SIGNAT	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 90/400mg Tablets □ 90/400mg Tablets □ 100/40mg Tablets □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day □ 1000mg per day □ 1200mg per day □ 1200mg per day □ 1200mg Tablets □ 550mg Tablets □ 550mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take 90mg daily with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one tablets orally o □ Take one 400mg tablet or Take two tablets once dailed to Take three tablets in the mith a meal, as directed our Take three tablets once dailed Take one tablet orally once Take 200mg tablet every ruled Take 400mg tablet every ruled Take 600mg tablet every ruled Take 600mg tablet every ruled Take 500mg tablet every ruled 500mg tablet every ruled 600mg	Direction ithout food in combination with Sovaldi® ithout food in combination with Sovaldi® ithout food in combination with Sovaldi® or without food or without food or without food orally once a day ally once a day ly in the morning with a meal morning and one tablet in the evening on the daily dose pack aily with food e daily with food morning/400mg tablet every evening morning/400mg tablet every evening morning/400mg tablet every evening morning/600mg tablet every evening morning/600mg tablet every evening worning/600mg tablet every evening morning/600mg tablet overy evening worning/600mg tablet overy evening	28 28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28 28 30 4 Carton 28 4 Carton 28 4 Carton 28 5 Carton 28 6 Carton 20 7 Car
Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO TM □ SOVALDI® □ TECHNIVIE TM □ VIEKIRA PAK TM □ VIEKIRA XR TM □ VOSEVI TM □ MODERIBA Dose Pack TM □ RIBASPHERE RibaPack® □ MODERIBA TM □ RIBASPHERE® □ RIBAVIRIN □ XIFAXAN® □ ZEPATIER TM □ U	□ 30mg Tablets □ 60mg Tablets □ 90mg Tablets □ 90/400mg Tablets □ 90/400mg Tablets □ 100/40mg Tablets □ 150mg Capsules □ 400mg Tablets □ 12.5/75/50mg Tablets □ 12.5/75/50mg & 250mg Dose Pack □ 200/8.33/50/33.33mg Tablets □ 400/100/100mg Tablets □ 600mg per day □ 1000mg per day □ 1200mg per day □ 1200mg per day □ 1200mg Tablets □ 550mg Tablets □ 550mg Tablets □ 50/100mg Tablets	□ Take 30mg daily with or w with or without Ribavirin □ Take 60mg daily with or w with or without Ribavirin □ Take 90mg daily with or w with or without Ribavirin □ Take 90mg daily with or without Ribavirin □ Take one tablet daily with □ Take one tablet daily with □ Take one tablets orally o □ Take one 400mg tablet or Take two tablets once dailed to Take three tablets in the mith a meal, as directed our Take three tablets once dailed Take one tablet orally once Take 200mg tablet every ruled Take 400mg tablet every ruled Take 600mg tablet every ruled Take 600mg tablet every ruled Take 500mg tablet every ruled 500mg tablet every ruled 600mg	Direction //ithout food in combination with Sovaldi® or without food or without food ore daily with food orally once a day ally once a day ly in the morning with a meal morning and one tablet in the evening or the daily dose pack aily with food morning/400mg tablet every evening morning/400mg tablet every evening morning/400mg tablet every evening morning/600mg tablet every evening //capsules every morning and, //capsules every evening y with or without food or without food	28 28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28