

HEPATITIS C VIRUS SPECIALTY CARE PROGRAM

Phone: **844-422-6400** • Fax: **888-850-4018**



v9.6_012217

©2017 KloudScript, Inc. - All rights reserved.

PATIENT INFORMATI		2 PRESCRIBER INFORMATION:		
Name:				
Address:		Address:		
City:				
Phone:				
Email:				
DOB: Gender: O				
Height: Weight:	Allergies:	Office Contact: Phone:		
STATEMENT OF MEI	DICAL NECESSITY: (Plea	ase Attach All Medical Documentation)		
	Diagnostic Information	Labs		
Date of Diagnosis: ICD-	-10: Race:	ALT: HGB:		
Genotype: Subtype: _	type: Subtype: Q80K: Desitive Degative (For Genotype 1a)			
Indicate Patient Status: ☐ Naïve ☐ Partial Responder ☐ Non-responder ☐ Null-responder ☐ Indicate Patient Status: ☐ Naïve ☐ Partial Responder ☐ Non-responder ☐ Null-responder ☐ Indicate Patient Status: ☐ Naïve ☐ Partial Responder ☐ Non-responder ☐ Null-responder ☐ Indicate Patient Status: ☐ Naïve ☐ Partial Responder ☐ Non-responder ☐ Null-responder ☐ Indicate Patient Status: ☐ Naïve ☐ Partial Responder ☐ Non-responder ☐ Null-responder ☐ Indicate Patient Status: ☐ Null-responder ☐ Indicate Patient Status: ☐ Null-responder ☐ Null-responder ☐ Null-responder ☐ Indicate Patient Status: ☐ Null-responder ☐ Null-respon		I Null-responder La Relapser		
Cirrhosis: ☐ No ☐ Yes If Yes: ☐ Compensated ☐ Decompensated History of Liver Biopsy? ☐ No ☐ Yes If Yes, Please Attach Results		NS5A Resistance Assay:	Date:	
☐ Fibrosure or ☐ Fibroscan: Results:		Medication List and Contra	aindications	
Extra-Hepatic Manifestations: Ascites Hepatic Encephalopathy Thrombocytopenia				
Other:		antation? Yes No		
HBsAg and anti-HBc Test: ☐ Positive ☐ Negative Date:				
If Prior Authorization is Denied:		☐ Anxiety ☐ Depression ☐ Pulmona	,	
☐ Automatically Draft Appeal for Re	view Send Preferred Formulary	Alternatives Renal Insufficiency Other:		
1 INJECTION TRAINING	G: O To Be Administered by Pharmacist	t O Pharmacist to Provide Training O Patient Trained in MD Office O Manu	ıfacturer Nurse Support	
5 PICK UP OR DELIVE	RY: O Delivery to Patient's	Home O Delivery to Physician's Office O Pharmacy	to Coordinate	
PRESCRIPTION: Duration of	of Therany: □ 8 Weeks □ 12 W	lacks D 21 Weeks D Other		
		Patient's Date of Birth:		
	Dosage & Strength	Patient's Date of Birth: Direction	QTY Refills	
Patient Name:		Patient's Date of Birth: Direction □ Take 30mg daily with or without food in combination with Sovaldi®	QTY Refills	
Patient Name:	Dosage & Strength	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi®		
Patient Name: Medication	Dosage & Strength ☐ 30mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin	28	
Patient Name: Medication	Dosage & Strength 30mg Tablets 60mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi®	28	
Patient Name: Medication □ DAKLINZA [™]	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin	28 28 28	
Patient Name: Medication □ DAKLINZA [™] □ EPCLUSA®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 400/100mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food	28 28 28 28	
Patient Name: Medication □ DAKLINZA [™] □ EPCLUSA® □ HARVONI® □ MAVYRET [™] □ OLYSIO®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90mg Tablets 100/400mg Tablets 100/40mg Tablets 150mg Capsules	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take three tablets orally once daily with food Take one 150mg capsule orally once a day	28 28 28 28 28 28 1 Carton 28	
Patient Name: Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO® □ SOVALDI®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90mg Tablets 100/400mg Tablets 100/40mg Tablets 150mg Capsules 400mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take three tablets orally once daily with food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day	28 28 28 28 28 28 1 Carton 28 28	
Patient Name: Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO® □ SOVALDI® □ TECHNIVIE TM	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90mg Tablets 100/40mg Tablets 150mg Capsules 400mg Tablets 12.5/75/50mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take three tablets orally once daily with food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal	28 28 28 28 28 28 1 Carton 28 28 28	
Patient Name: Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO® □ SOVALDI®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90mg Tablets 100/400mg Tablets 100/400mg Tablets 150mg Capsules 400mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg & 250mg	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take three tablets orally once daily with food Take one 150mg capsule orally once a day Take two tablets once daily in the morning with a meal	28 28 28 28 28 28 1 Carton 28 28	
Patient Name: Medication □ DAKLINZA TM □ EPCLUSA® □ HARVONI® □ MAVYRET TM □ OLYSIO® □ SOVALDI® □ TECHNIVIE TM	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90mg Tablets 100/40mg Tablets 150mg Capsules 400mg Tablets 12.5/75/50mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take three tablets orally once daily with food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal	28 28 28 28 28 28 1 Carton 28 28 28	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 400/100mg Tablets 90/400mg Tablets 100/40mg Tablet 150mg Capsules 400mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg & 250mg Dose Pack	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take three tablets orally once daily with food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal Take three tablets in the morning and one tablet in the evening with a meal, as directed on the daily dose pack	28 28 28 28 28 28 1 Carton 28 28 56 1 Pack	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM VOSEVI®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90mg Tablets 100/40mg Tablets 100/40mg Tablets 150mg Capsules 12.5/75/50mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg & 250mg Dose Pack 200/8.33/50/33.33mg Tablets 400/100/100mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal Take three tablets in the morning and one tablet in the evening with a meal, as directed on the daily dose pack Take one tablet orally once daily with food Take one tablet orally once daily with food Take 200mg tablet every morning/400mg tablet every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM VIEKIRA XR TM VOSEVI® MODERIBA Dose Pack TM	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90/400mg Tablets 100/40mg Tablets 1150mg Capsules 400mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg Tablets 400/100/100mg Tablets 600mg per day 800mg per day	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take three tablets orally once daily with food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal Take three tablets in the morning and one tablet in the evening with a meal, as directed on the daily dose pack Take three tablets once daily with food Take one tablet orally once daily with food Take 200mg tablet every morning/400mg tablet every evening Take 400mg tablet every morning/400mg tablet every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM VOSEVI®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90mg Tablets 100/40mg Tablets 100/40mg Tablets 150mg Capsules 12.5/75/50mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg & 250mg Dose Pack 200/8.33/50/33.33mg Tablets 400/100/100mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal Take three tablets in the morning and one tablet in the evening with a meal, as directed on the daily dose pack Take one tablet orally once daily with food Take one tablet orally once daily with food Take 200mg tablet every morning/400mg tablet every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM VIEKIRA XR TM VOSEVI® MODERIBA Dose Pack TM RIBASPHERE RibaPack®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90/400mg Tablets 90/400mg Tablets 100/40mg Tablets 150mg Capsules 400mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg Tablets 400/100/100mg Tablets 600mg per day 800mg per day 1000mg per day	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take three tablets orally once daily with food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal Take three tablets in the morning and one tablet in the evening with a meal, as directed on the daily dose pack Take three tablets once daily with food Take one tablet orally once daily with food Take 200mg tablet every morning/400mg tablet every evening Take 400mg tablet every morning/400mg tablet every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM VIEKIRA XR TM VOSEVI® MODERIBA Dose Pack TM RIBASPHERE RibaPack®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90/400mg Tablets 90/400mg Tablets 100/40mg Tablets 150mg Capsules 400mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg Tablets 200/8.33/50/33.33mg Tablets 400/100/100mg Tablets 600mg per day 800mg per day 1000mg per day 1200mg per day 1200mg per day 200mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take one tablet daily with or without food Take one tablets orally once daily with food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal Take three tablets in the morning and one tablet in the evening with a meal, as directed on the daily dose pack Take three tablets once daily with food Take one tablet orally once daily with food Take 200mg tablet every morning/400mg tablet every evening Take 600mg tablet every morning/400mg tablet every evening Take 600mg tablet every morning/600mg tablet every evening Take tablets/capsules every morning and, Take tablets/capsules every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM VIEKIRA YAK TM VOSEVI® MODERIBA Dose Pack TM RIBASPHERE RibaPack® MODERIBA TM RIBASPHERE® RIBAVIRIN XIFAXAN®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90/400mg Tablets 100/40mg Tablets 1100/40mg Tablets 120mg Capsules 12.5/75/50mg Tablets 12.5/75/50mg & 250mg Dose Pack 200/8.33/50/33.33mg Tablets 400/100/100mg Tablets 600mg per day 800mg per day 1000mg per day 1200mg per day 1200mg Tablets 200mg Tablets	Patient's Date of Birth: Direction	28 28 28 28 28 1 Carton 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM VIEKIRA YAK TM VOSEVI® MODERIBA Dose Pack TM RIBASPHERE RibaPack® MODERIBA TM RIBASPHERE® RIBAVIRIN XIFAXAN® ZEPATIER®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90/400mg Tablets 90/400mg Tablets 100/40mg Tablets 150mg Capsules 400mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg Tablets 200/8.33/50/33.33mg Tablets 400/100/100mg Tablets 600mg per day 800mg per day 1000mg per day 1200mg per day 1200mg per day 200mg Tablets	Patient's Date of Birth: Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take one tablet daily with or without food Take one tablets orally once daily with food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal Take three tablets in the morning and one tablet in the evening with a meal, as directed on the daily dose pack Take three tablets once daily with food Take one tablet orally once daily with food Take 200mg tablet every morning/400mg tablet every evening Take 600mg tablet every morning/400mg tablet every evening Take 600mg tablet every morning/600mg tablet every evening Take tablets/capsules every morning and, Take tablets/capsules every evening	28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM VIEKIRA XR TM VOSEVI® MODERIBA Dose Pack TM RIBASPHERE RibaPack® MODERIBA TM RIBASPHERE® RIBAVIRIN XIFAXAN® ZEPATIER®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90/400mg Tablets 100/40mg Tablets 1150mg Capsules 12.5/75/50mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg & 250mg Dose Pack 200/8.33/50/33.33mg Tablets 400/100/100mg Tablets 600mg per day 800mg per day 11000mg per day 1200mg per day 1200mg per day 1200mg per day 1200mg Capsules 550mg Tablets 550/100mg Tablets	Direction □ Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin □ Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin □ Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin □ Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin □ Take one tablet daily with or without food □ Take one tablet daily with or without food □ Take one 150mg capsule orally once a day □ Take one 150mg capsule orally once a day □ Take two tablets once daily in the morning with a meal □ Take three tablets in the morning and one tablet in the evening with a meal, as directed on the daily dose pack □ Take three tablets once daily with food □ Take one tablet orally once daily with food □ Take 200mg tablet every morning/400mg tablet every evening □ Take 400mg tablet every morning/400mg tablet every evening □ Take 600mg tablet every morning/400mg tablet every evening □ Take 600mg tablet every morning/600mg tablet every evening □ Take one tablet twice daily with or without food □ Take one tablet twice daily with or without food	28 28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28	
Patient Name: Medication DAKLINZA TM EPCLUSA® HARVONI® MAVYRET TM OLYSIO® SOVALDI® TECHNIVIE TM VIEKIRA PAK TM VIEKIRA XR TM VOSEVI® MODERIBA Dose Pack TM RIBASPHERE RibaPack® RIBASPHERE® RIBAVIRIN XIFAXAN® ZEPATIER®	Dosage & Strength 30mg Tablets 60mg Tablets 90mg Tablets 90/400mg Tablets 100/40mg Tablets 1100/40mg Tablets 112.5/75/50mg Tablets 12.5/75/50mg Tablets 12.5/75/50mg & 250mg Dose Pack 200/8.33/50/33.33mg Tablets 600mg per day 800mg per day 1000mg per day 1200mg per day 1200mg per day 1200mg Tablets 550mg Tablets 550mg Tablets	Direction Take 30mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 60mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin Take one tablet daily with or without food Take one tablet daily with or without food Take one 150mg capsule orally once a day Take one 400mg tablet orally once a day Take two tablets once daily in the morning with a meal Take three tablets in the morning and one tablet in the evening with a meal, as directed on the daily dose pack Take three tablets once daily with food Take one tablet orally once daily with food Take 200mg tablet every morning/400mg tablet every evening Take 400mg tablet every morning/400mg tablet every evening Take 600mg tablet every morning/400mg tablet every evening Take 600mg tablet every morning/600mg tablet every evening Take tablets/capsules every morning and, Take tablets/capsules every evening Take one tablet twice daily with or without food Take one tablet daily with or without food	28 28 28 28 28 28 1 Carton 28 28 56 1 Pack 1 Carton 28	