

## INFLAMMATORY BOWEL DISEASE SPECIALTY CARE PROGRAM

Phone: 844-422-6400 •

Fax: 888-850-4018



v9.0\_032917

©2017 KloudScript, Inc. - All rights reserved.

## PATIENT INFORMATION:

## PRESCRIBER INFORMATION:

Name:	Name:
Address:	Address:
City: State: Zip:	City: State: Zip:
Phone: Alt. Phone:	Phone: Fax:
Email:	NPI: DEA:
DOB: Gender: O M O F Caregiver:	Tax I.D.:
Height: Weight: Allergies:	Office Contact: Phone:

## 3 STATEMENT OF MEDICAL NECESSITY: (Please Attach All Medical Documentation)

Date of Diagnosis:	Prior	Indicate Drug Name
□ Crohn's Disease □ Ulcerative Colitis □ Irritable Bowe		s: and Length of Treatment:
ICD-10:	□ 5-ASA	
Other:	Biologics	
Serious or active infection present?	Corticosteroids	
Hep B ruled out or treatment started?	□ Immunosuppre	ssants
TB Test: Dositive Negative Date:	Methotrexate	
If Prior Authorization is Denied:		
Automatically Draft Appeal for Review		
Send Preferred Formulary Alternatives	🗖 Other	

**1** INJECTION TRAINING: O To Be Administered by Pharmacist O Pharmacist to Provide Training O Patient Trained in MD Office O Manufacturer Nurse Support

**5 PICK UP OR DELIVERY:** O Delivery to Patient's Home O Delivery to Physician's Office O Pharmacy to Coordinate

6 INSURANCE INFORMATION: Please Include Front and Back Copies of Pharmacy and Medical Card

**PRESCRIPTION INFORMATION:** (Please be sure to choose both induction and maintenance dose where applicable)

Patient Name:

Patient's Date of Birth:

Medication	Dosage & Strength	Direction	QTY	Refills	
	Prefilled Syringe Starter Kit	□ Induction Dose: Inject 400mg SC on day 1, 14 and 28	6	0	
	<ul> <li>200mg/ml Prefilled Syringe</li> <li>200mg Lyophilized Powder</li> </ul>	Maintenance: Inject 400mg SC every 4 weeks	2		
	Crohn's Disease/ Ulcerative Colitis Starter Kit	Induction Dose: Inject 160mg SC on day 1, then 80mg SC on day 15, then switch to maintenance dose	6 0		
	<ul> <li>40mg/0.8ml Pen</li> <li>40mg/0.8ml Prefilled Syringe</li> </ul>	Maintenance: Inject 40mg SC every other week	2		
Patient has signed HUMIRA Complete form					
	100mg/ml Smartject <sup>®</sup> Autoinjector	Induction Dose: Inject 200mg SC at week 0, 100mg SC at week 2 and then switch to maintenance dose	3	0	
	100mg/ml Prefilled Syringe	Maintenance: Inject 100mg SC every 4 weeks	1		
	□ 130mg/26ml Vial	Induction Dose: Patient Weight <55kg: 260mg; >55kg to 85kg: 390mg; >85 kg: 520mg administered IV		0	
	<ul> <li>45mg/0.5ml Prefilled Syringe</li> <li>90mg/ml Prefilled Syringe</li> <li>45mg/0.5ml Vial</li> </ul>	□ Maintenance Dose: Inject 90mg SC 8 weeks after the initial intravenous dose, then every 8 weeks thereafter	1		
	9mg Tablets	Take one tablet daily in the morning with or without food	30	1	
	550mg Tablets	Take one tablet three times daily for 14 days	42		
•					
PRESCRIBER SIGNATURE:   authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs.					
		Signature: E			
Substitution Permitted Dispense As Written Prior authorization approval and insurance benefits will be determined by the payor based upon the patient's eligibility, medical necessity, and the terms of the patient's coverage, among other things. Participation in this program is not a guarantee of prior authorization or of payment.					

Confidentiality Notice: This fax is intended to be delivered only to the named addressee and contains confidential information that may be protected health information under federal and state laws. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please inform the sender immediately if you have received this document in error and then destroy this document immediately.