

RHEUMATOID ARTHRITIS SPECIALTY CARE PROGRAM

Phone: 844-422-6400 Fax: 888-850-4018



ame:		Name:		
aaress:	State: Zip:	Address: State:	7in.	
	State: Zip: Alt. Phone:			
mail:	Alt. Filone	Priorie: Pax NPI: DEA:		
Παιι ΩΒ·	ender: O M O F Caregiver:	NFI BEA		
eight: We	eight: Allergies:	Office Contact: Phone:		
			nents:	
	OF MEDICAL NECESSITY: (Please Att	dach All Medical Documentation) Azulfidine® Celebrex®	☐ Me	ethotrexa
ate of Diagnosis:	Patient also taking Metho	otrexate?	ds 🔟 Ot	hers
DD-10:	Serious or active infectior Hep B ruled out or treatm	Indicate Drug Name and Lend	gth of Trea	itment:
other:	Hep B ruled out or treatm Does patient have latex a	llergy? ☐ Yes ☐ No		
B Test: Desitive Desi	egative Date: LFT: ALT: AST:	Date:		
f Prior Authorization is	denied, recommended formulary alternatives	will be provided to the prescriber based upon the patient's in	surance c	overage
INJECTION T	RAINING: O To Be Administered by Pharmacist	O Pharmacist to Provide Training O Patient Trained in MD Office O Mar	nufacturer Nu	rse Suppor
PICK UP OR I	DELIVERY: O Delivery to Patient's Ho	ome O Delivery to Physician's Office O Pharmacy	to Coord	dinate
INSURANCE I	NFORMATION: Please Include Front	and Back Copies of Pharmacy and Medical Card		
	INFORMATION: (Please be sure to c	choose both induction and maintenance dose wher		
atient Name: Medication	Dosage & Strength	Patient's Date of Birth: Direction	QTY	
Medication	Dosage & Strength	☐ Inject 162mg SC every other week (< 220 lbs)	QII	Reilli
☐ ACTEMRA®	☐ 162mg/0.9ml Prefilled Syringe	☐ Inject 162mg SC every week (> 220 lbs) ☐ Inject 162mg SC every 2 weeks (> 66lbs) ☐ Inject 162mg SC every 3 weeks (< 66lbs)		
□ CIMZIA®	☐ Prefilled Syringe Starter Kit☐ 200mg/ml Prefilled Syringe☐ 200mg Lyophilized Powder Vial☐ Prefilled Syringe☐ 200mg Lyophilized Powder Vial☐ Prefilled Syringe	☐ Induction Dose: Inject 400mg SC on day 1, day 14 and day 28 ☐ Maintenance: Inject 400mg SC every 4 weeks ☐ Maintenance: Inject 200mg SC every other week	6 2	0
□ COSENTYX®	☐ 150mg/ml Sensoready® Pen☐ 150mg/ml Prefilled Syringe☐ 150mg/ml Lyophilized Powder Vial	☐ Induction Dose: Inject 150mg SC at weeks 0, 1, 2, 3, and 4☐ Induction Dose: Inject 300mg SC at weeks 0, 1, 2, 3, and 4☐ Maintenance Dose: Inject 150mg SC every four weeks☐ Maintenance Dose: Inject 300mg SC every four weeks☐ Maintenance Dose: Inject 150mg SC every four weeks☐ Maintenance Dose every four weeks☐ Maintenance Dose	5 10 1 2	0
□ ENBREL®	□ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge □ For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/ml Vial □ 25mg/ml Vial	☐ Inject 50mg SC once a week ☐ Inject 25mg SC twice a week (3-4 days apart) ☐ Other	4 1	
☐ HUMIRA®	☐ 40mg/0.8ml Pen☐ 40mg/0.8ml Prefilled Syringe	☐ Inject 40mg SC every other week ☐ Patient has signed ☐ Inject 40mg SC once a week ☐ HUMIRA Complete form		
□ KEVZARA®	☐ 150mg/1.14ml Prefilled Syringe	☐ Inject 150mg SC every 2 weeks	2	
U NEVZANA*	□ 200mg/1.14ml Prefilled Syringe	☐ Inject 200mg SC every 2 weeks ☐ Induction Dose: Patient Weight < 132 lbs: 500mg; 132-220 lbs:	2	
□ ORENCIA®	☐ 250mg Lyophilized Powder Vial ☐ 125mg/M ClickJect™ Autoinjector	750mg; > 220 lbs: 1000mg administered IV, then inject 125mg SC within 24 hours		0
	 □ 50mg/0.4ml Prefilled Syringe □ 87.5mg/0.7ml Prefilled Syringe □ 125mg/ml Prefilled Syringe 	☐ Inject 50mg SC once a week (10 to less than 25kg)☐ Inject 87.5mg SC once a week (25 to less than 50kg)☐ Inject 87.5mg SC once a week (2	4	
OTEZLA® (for PsA)	□ Starter Pack (Titration) □ 30mg Tablets	☐ Inject 125mg SC once a week (50kg or more) ☐ Starter Pack: Take one tablet in the morning on day 1, then take one tablet in the morning and one tablet in the evening as	1	0
		directed on the starter pack		
	☐ 50mg/0.5ml Smartject Autoinjector	☐ Maintenance: Take one 30mg tablet by mouth twice daily ☐ Inject 50mg SC once a month	60	+
□ SIMPONI®	□ 50mg/0.5ml Prefilled Syringe □ 45mg/0.5ml Prefilled Syringe (for < 220 lbs)	☐ Induction Dose: Inject 1 prefilled syringe SC on day 1	1	
STELARA® (for PsA)	☐ 90mg/1ml Prefilled Syringe (for > 220 lbs)	☐ Maintenance: Inject 1 prefilled syringe SC on day 29, and every 12 weeks thereafter	1	0
(.5 5/ 9	☐ Yes or ☐ No: STELARA SELF-INJECTION: Healthcare provider certifies	s that patient has been trained and is eligible for self-injection		<u> </u>
☐ TALTZ®	□ 80mg/ml Single-Dose Prefilled Autoinjector□ 80mg/ml Single-Dose Prefilled Syringe	☐ Induction Dose: Inject 160mg SC (two 80mg injections) at weeks 0☐ Maintenance: Inject 80mg SC every 4 weeks	2	0
□ XELJANZ®	□ 5mg Tablet	☐ Take one 5mg tablet by mouth twice a day	60	+
☐ XELJANZ® XR	☐ 11mg Tablet	☐ Take one 11mg tablet once a day	30	
	<u> </u>	<u> </u>		
□ RASUVO®	_		1	
		nee for initiating and coordinating insurance prior authorizations, nursing services and pat	tiont ancietes	o progress: